



INDICATIONS

ISENTRESS is an anti-HIV medicine used for the treatment of HIV. ISENTRESS must be used with other anti-HIV medicines, which may increase the likelihood of response to treatment.

The safety and effectiveness of ISENTRESS in children has not been studied.

It is important that you remain under your doctor's care.

ISENTRESS will NOT cure HIV infection or reduce your chance of passing HIV to others through sexual contact, sharing needles, or being exposed to your blood.

IMPORTANT RISK INFORMATION

A condition called Immune Reconstitution Syndrome can happen in some patients with advanced HIV infection (AIDS) when anti-HIV treatment is started. Signs and symptoms of inflammation from opportunistic infections may occur as the medicines work to treat the HIV infection and strengthen the immune system. Call your doctor right away if you notice any signs or symptoms of an infection after starting ISENTRESS.

Contact your doctor immediately if you experience unexplained muscle pain, tenderness, or weakness while taking ISENTRESS. This is because on rare occasions muscle problems can be serious and can lead to kidney damage.

When ISENTRESS has been given with other anti-HIV drugs, side effects included nausea, headache, tiredness, weakness, trouble sleeping, stomach pain, dizziness, depression, and suicidal thoughts and actions.

Rash occurred more often in patients taking ISENTRESS and darunavir together than with either drug separately, but was generally mild.