

- KALETRA oral solution that is stored at room temperature (less than 77°F or 25°C) should be used within 2 months.
 - Keep KALETRA away from high heat.
- Throw away any medicine that is out of date or that you no longer need.

Keep KALETRA and all medicines out of the reach of children.

General information about KALETRA

KALETRA does not cure HIV-1 or AIDS. The long-term effects of KALETRA are not known at this time. People taking KALETRA may still get opportunistic infections or other conditions that happen with HIV-1 infection. Some of these conditions are pneumonia, herpes virus infections, and *Mycobacterium avium* complex (MAC) infections.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use KALETRA for a condition for which it was not prescribed. Do not give KALETRA to other people, even if they have the same condition you have. It may harm them.

This Medication Guide summarizes the most important information about KALETRA. If you would like more information, talk with your doctor. You can ask your pharmacist or doctor for information about KALETRA that is written for health professionals. For more information about KALETRA call 1-800-633-9110 or go to www.KALETRA.com.

What are the ingredients in KALETRA?

Active ingredient: lopinavir and ritonavir
Inactive ingredients:

KALETRA 200 mg lopinavir and 50 mg ritonavir tablets: copovidone, sorbitan monolaurate, colloidal silicon dioxide, and sodium stearyl fumarate. The film coating contains: hypromellose, titanium dioxide, polyethylene glycol 400, hydroxypropyl cellulose, talc, colloidal silicon dioxide, polyethylene glycol 3350, yellow ferric oxide 172, and polysorbate 80.

KALETRA 100 mg lopinavir and 25 mg ritonavir tablets: copovidone, sorbitan monolaurate, colloidal silicon dioxide, and sodium stearyl fumarate. The film coating contains: polyvinyl alcohol, titanium dioxide, talc, polyethylene glycol 3350, and yellow ferric oxide E172.

KALETRA oral solution: acesulfame potassium, alcohol, artificial cotton candy flavor, citric acid, glycerin, high fructose corn syrup, Magnasweet-110 flavor, menthol, natural and artificial vanilla flavor, peppermint oil, polyoxyl 40 hydrogenated castor oil, povidone, propylene glycol, saccharin sodium, sodium chloride, sodium citrate, and water.

KALETRA oral solution contains 42.4% alcohol (v/v). "See How should I take KALETRA?"

KALETRA Tablets, 200 mg lopinavir/50 mg ritonavir
Manufactured by Abbott Pharmaceuticals PR Ltd.,
Barceloneta, PR 00617
for Abbott Laboratories, North Chicago, IL 60064, U.S.A.
KALETRA Tablets, 100 mg lopinavir/25 mg ritonavir and
KALETRA Oral Solution
Abbott Laboratories, North Chicago, IL 60064, U.S.A.
2010, ALL RIGHTS RESERVED

* The brands listed are trademarks of their respective owners and are not trademarks of Abbott Laboratories. The makers of these brands are not affiliated with and do not endorse Abbott Laboratories or its products.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Ref: 03-A387-R8

Revised: June, 2010

036-395113 MASTER

039-418110

