



STALLION.

It is frequently the case that a tree which has received all the care and attention which can be bestowed upon it by the most experienced nurseryman, is transplanted to a soil of very inferior character, and being thus stunted in its growth, is the frequent cause of dissatisfaction to the purchaser.

REMARKS ON TRANSPLANTING TREES.

It is frequently the case that a tree which has received all the care and attention which can be bestowed upon it by the most experienced nurseryman, is transplanted to a soil of very inferior character, and being thus stunted in its growth, is the frequent cause of dissatisfaction to the purchaser.

Many of the most experienced cultivators regard the Fall, immediately after the first hard frost has arrested the growth, as the best season for transplanting every variety of trees but evergreens, which should be planted during the last days of Spring, or the first of summer.

When the tree has been some time out of the ground, it is well to immerse the bodies and roots in water for about twenty-four hours; this will much benefit the tree, and advance its vegetation.

When the hole is three quarters filled, pour in three or four gallons of water, and after it has settled away, fill up the hole, pressing the earth around the tree with the foot.

SOWING WINTER GRAIN.

What.—It is not always easy to tell whether early or late sowing would be most advisable. Early sown grain gets the best hold against the winter, and when it is not liable to attacks from insects, will generally yield best.

Unfortunately happens that the Hessian fly is usually more injurious to early than late sown grain because a generation of the insect is sometimes brought forward in the fall.

Preparation of the ground.—In this operation, regard should be had to protecting the crop from injury during winter.

Many of the most experienced cultivators regard the Fall, immediately after the first hard frost has arrested the growth, as the best season for transplanting every variety of trees but evergreens, which should be planted during the last days of Spring, or the first of summer.

When the tree has been some time out of the ground, it is well to immerse the bodies and roots in water for about twenty-four hours; this will much benefit the tree, and advance its vegetation.

When the hole is three quarters filled, pour in three or four gallons of water, and after it has settled away, fill up the hole, pressing the earth around the tree with the foot.

SOWING WINTER GRAIN.

What.—It is not always easy to tell whether early or late sowing would be most advisable. Early sown grain gets the best hold against the winter, and when it is not liable to attacks from insects, will generally yield best.

Unfortunately happens that the Hessian fly is usually more injurious to early than late sown grain because a generation of the insect is sometimes brought forward in the fall.

Preparation of the ground.—In this operation, regard should be had to protecting the crop from injury during winter.

Many of the most experienced cultivators regard the Fall, immediately after the first hard frost has arrested the growth, as the best season for transplanting every variety of trees but evergreens, which should be planted during the last days of Spring, or the first of summer.

When the tree has been some time out of the ground, it is well to immerse the bodies and roots in water for about twenty-four hours; this will much benefit the tree, and advance its vegetation.

When the hole is three quarters filled, pour in three or four gallons of water, and after it has settled away, fill up the hole, pressing the earth around the tree with the foot.

SOWING WINTER GRAIN.

What.—It is not always easy to tell whether early or late sowing would be most advisable. Early sown grain gets the best hold against the winter, and when it is not liable to attacks from insects, will generally yield best.

Unfortunately happens that the Hessian fly is usually more injurious to early than late sown grain because a generation of the insect is sometimes brought forward in the fall.

Preparation of the ground.—In this operation, regard should be had to protecting the crop from injury during winter.

Many of the most experienced cultivators regard the Fall, immediately after the first hard frost has arrested the growth, as the best season for transplanting every variety of trees but evergreens, which should be planted during the last days of Spring, or the first of summer.

When the tree has been some time out of the ground, it is well to immerse the bodies and roots in water for about twenty-four hours; this will much benefit the tree, and advance its vegetation.

When the hole is three quarters filled, pour in three or four gallons of water, and after it has settled away, fill up the hole, pressing the earth around the tree with the foot.

SOWING WINTER GRAIN.

What.—It is not always easy to tell whether early or late sowing would be most advisable. Early sown grain gets the best hold against the winter, and when it is not liable to attacks from insects, will generally yield best.

Unfortunately happens that the Hessian fly is usually more injurious to early than late sown grain because a generation of the insect is sometimes brought forward in the fall.

Preparation of the ground.—In this operation, regard should be had to protecting the crop from injury during winter.

Many of the most experienced cultivators regard the Fall, immediately after the first hard frost has arrested the growth, as the best season for transplanting every variety of trees but evergreens, which should be planted during the last days of Spring, or the first of summer.

When the tree has been some time out of the ground, it is well to immerse the bodies and roots in water for about twenty-four hours; this will much benefit the tree, and advance its vegetation.

When the hole is three quarters filled, pour in three or four gallons of water, and after it has settled away, fill up the hole, pressing the earth around the tree with the foot.

SOWING WINTER GRAIN.

What.—It is not always easy to tell whether early or late sowing would be most advisable. Early sown grain gets the best hold against the winter, and when it is not liable to attacks from insects, will generally yield best.

Unfortunately happens that the Hessian fly is usually more injurious to early than late sown grain because a generation of the insect is sometimes brought forward in the fall.

Preparation of the ground.—In this operation, regard should be had to protecting the crop from injury during winter.

Many of the most experienced cultivators regard the Fall, immediately after the first hard frost has arrested the growth, as the best season for transplanting every variety of trees but evergreens, which should be planted during the last days of Spring, or the first of summer.

When the tree has been some time out of the ground, it is well to immerse the bodies and roots in water for about twenty-four hours; this will much benefit the tree, and advance its vegetation.

When the hole is three quarters filled, pour in three or four gallons of water, and after it has settled away, fill up the hole, pressing the earth around the tree with the foot.

SOWING WINTER GRAIN.

What.—It is not always easy to tell whether early or late sowing would be most advisable. Early sown grain gets the best hold against the winter, and when it is not liable to attacks from insects, will generally yield best.

PARSONS & CO. The silk culture is rapidly extending in Tennessee. Large quantities of cocoons are raised and sold at \$3 per bushel, and a company with a capital of \$100,000, is located on the Red River, 35 miles below Nashville, engaged in the exclusive manufacture of silks.