

## Communication

To The Taxpayers  
Of The City Of Landrum

and especially the over 400 signatures who had the intelligence and foresight to put their names on the dotted line to demand an Ordinance for the control of the dog population, which we all know has gotten out of hand. Recently a lady has been bitten by a dog. This is a serious matter. Complaints of dog harassment have been rudely ignored in the past. A crucial meeting on this matter will take place on Saturday, March 22nd at 10:00 A.M. at City Hall. We shall find how progressive our City Government really is. All those who want to stop this backwoods tradition to allow dogs to run loose, to manure your front lawns, porches and yards, destroy your shubbery and gardens, harass your children and pets, attack adults and livestock, bark at all hours and keep you awake, spread disease to other animals and humans, please attend this meeting and assert yourselves in no uncertain terms. In my past attempts to convince Council that my proposals are realistic, that it will separate the wanted cared for pets from the unwanted stray dogs, produced no encouraging results. My proposals were either only glanced over, misunderstood, blown out of context, ridiculed, soft peddled and no intelligent counter proposal made.

At the last Council meeting, Jack Nodine committed a serious blunder by insulting the public with the outrageous remark, "If you don't like dogs running loose move out of town."

It is urged that all interested citizens of Landrum be present at the Saturday, March 22nd meeting.

Edmund A. Komant  
Chairman  
Dog Ordinance Committee

The Hawaiian Islands grew from a chain of volcanoes. Oahu is the oldest of the islands, and none of its volcanoes is active. Diamond Head, the dramatic landmark that towers over Waikiki Beach, is a wall of the huge crater of one such extinct volcano, says National Geographic World magazine.

In foal thoroughbred mare for sale, \$1000. Call 894-8455 after 6 p.m. adv. 17-21,c.

WILL DO SEWING  
By Appointment Only  
Call Margaret Durham  
894-8785

adv. 19, 26, apr. 2, pd.

FOR SALE: Nice 3 bedroom mobile home, fully furnished with 23,000 BTU air conditioner. Will deliver \$4500. Call 859-5334. adv. 19, 20, 21, pd.

FOR SALE: 1973 350 Honda Good condition, \$325.00 or Triumph Chopper with a lot of extras \$925.00. Call 859-5334. adv. 19, 20, 21, pd.

# NATIONAL POISON PREVENTION WEEK

(March 16 thru March 25)

**AMERICAN DRUGGIST**  
**COUNTERDOSES FOR THE HOME**

<b>POISONS</b>  Acids • 16 After-Shave Lotion • 8 Camphor • 1 Carbon Monoxide • 11 Chlorine Bleach • 15 Cologne, Perfume • 8 Detergents • 15 Disinfectant with chlorine • 15 with carbolic acid • 3 Food Poisoning • 6 Furniture Polish • 14 Gasoline, Kerosene • 14 Household Ammonia • 13 Insect & Rat Poisons with arsenic • 2 with sodium fluoride • 10 with phosphorus • 12 with strychnine • 5 Lighter Fluid • 14 Lye • 13 Wild Mushrooms • 6 Oil of Wintergreen • 8 Pine Oil • 14 Rubbing Alcohol • 8 Turpentine • 14	<b>DO THIS FIRST</b>  <ul style="list-style-type: none"> <li>• CALL A PHYSICIAN—IMMEDIATELY!</li> <li>• Keep the patient warm.</li> <li>• Determine if the patient has taken                         <ul style="list-style-type: none"> <li>(1) A POISON</li> <li>(2) AN OVERDOSE</li> </ul> </li> <li>• While waiting for physician, give appropriate counterdose below.</li> <li>• But do not force any liquids on the patient—if he is unconscious.</li> </ul> <p style="font-size: small; margin-top: 5px;">• And do not induce vomiting if patient is having convulsions, or if patient is unconscious.</p> <p><b>To Find The Correct Counterdose</b></p> <ul style="list-style-type: none"> <li>• In one of the lists printed at left, find substance causing the trouble.</li> <li>• Next to that substance is a number. This refers to counterdose bearing same number in the section below.</li> </ul>	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; padding: 5px;"> <b>Keep All Poisons And Medicines Out Of Reach Of Children!</b> </td> <td style="width: 50%; padding: 5px;"> <b>1</b> Induce vomiting with                              • Finger in throat. OR                              • 1 tablespoon of syrup of ipecac, followed by a glass of water. OR                              • Teaspoonful of mustard in half glass of water                         </td> </tr> <tr> <td style="padding: 5px;"> <b>3</b> • Induce vomiting (See #1)                              • Then give 4 tablespoons of castor oil                              • Next give glass of milk OR the white of 2 raw eggs                         </td> <td style="padding: 5px;"> <b>4</b> • Induce vomiting (see #1) if patient is conscious.                              • Give glass of milk. OR activated charcoal in water                              • Give 2 tablespoons of epsom salt in 2 glasses of water.                              • Keep patient awake                         </td> </tr> <tr> <td style="padding: 5px;"> <b>6</b> • Induce vomiting (See #1)                              • Next give 2 tablespoons of epsom salt in 2 glasses of water—except in cases where diarrhea is severe                         </td> <td style="padding: 5px;"> <b>7</b> • Induce vomiting (See #1)                              • Give 2 teaspoons of bicarbonate of soda in a glass of warm water                              • Finally give glass of milk.                         </td> </tr> <tr> <td style="padding: 5px;"> <b>9</b> • Give activated charcoal in water.                              • Induce vomiting (see #1) but not with ipecac.                              • Give 2 tablespoons of epsom salt in 2 glasses of water.                         </td> <td style="padding: 5px;"> <b>10</b> • Give glass of milk or lime water.                              • Then induce vomiting. (See #1)                         </td> </tr> <tr> <td style="padding: 5px;"> <b>12</b> • Induce vomiting (See #1).                              • Don't let vomit touch victim's skin or yours.                              • Then give 4 oz mineral oil. Positively do NOT give vegetable or animal oil.                              • Also give 1 tablespoon of bicarbonate of soda in a quart of warm water                         </td> <td style="padding: 5px;"> <b>13</b> • Give 2 tablespoons of vinegar in 2 glasses of water                              • Then give the white of 2 raw eggs... or 2 ounces of vegetable oil.                              • Do NOT induce vomiting!                         </td> </tr> <tr> <td style="padding: 5px;"> <b>15</b> • Give patient one or two glasses of milk                         </td> <td style="padding: 5px;"> <b>16</b> • Give large quantity of water                              • Give 2 tablespoons of milk of magnesia, or a similar antacid.                              • Do NOT induce vomiting!                         </td> </tr> </table>	<b>Keep All Poisons And Medicines Out Of Reach Of Children!</b>	<b>1</b> Induce vomiting with • Finger in throat. OR • 1 tablespoon of syrup of ipecac, followed by a glass of water. OR • Teaspoonful of mustard in half glass of water	<b>3</b> • Induce vomiting (See #1) • Then give 4 tablespoons of castor oil • Next give glass of milk OR the white of 2 raw eggs	<b>4</b> • Induce vomiting (see #1) if patient is conscious. • Give glass of milk. OR activated charcoal in water • Give 2 tablespoons of epsom salt in 2 glasses of water. • Keep patient awake	<b>6</b> • Induce vomiting (See #1) • Next give 2 tablespoons of epsom salt in 2 glasses of water—except in cases where diarrhea is severe	<b>7</b> • Induce vomiting (See #1) • Give 2 teaspoons of bicarbonate of soda in a glass of warm water • Finally give glass of milk.	<b>9</b> • Give activated charcoal in water. • Induce vomiting (see #1) but not with ipecac. • Give 2 tablespoons of epsom salt in 2 glasses of water.	<b>10</b> • Give glass of milk or lime water. • Then induce vomiting. (See #1)	<b>12</b> • Induce vomiting (See #1). • Don't let vomit touch victim's skin or yours. • Then give 4 oz mineral oil. Positively do NOT give vegetable or animal oil. • Also give 1 tablespoon of bicarbonate of soda in a quart of warm water	<b>13</b> • Give 2 tablespoons of vinegar in 2 glasses of water • Then give the white of 2 raw eggs... or 2 ounces of vegetable oil. • Do NOT induce vomiting!	<b>15</b> • Give patient one or two glasses of milk	<b>16</b> • Give large quantity of water • Give 2 tablespoons of milk of magnesia, or a similar antacid. • Do NOT induce vomiting!	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="padding: 5px;"> <b>2</b> • Give glass of milk. OR 1 tablespoonful of activated charcoal mixed with a little water.                              • Induce vomiting (see #1) but not with syrup of ipecac if charcoal was given                         </td> </tr> <tr> <td style="padding: 5px;"> <b>5</b> • Give glass of milk, or activated charcoal in water.                              • Induce vomiting (#1) if not in convulsions. Do not use ipecac if charcoal was given.                              • Keep patient quiet.                         </td> </tr> <tr> <td style="padding: 5px;"> <b>8</b> • Give a glass of milk                              • Next induce vomiting (#1)                              • Give 2 teaspoons of bicarbonate of soda in a glass of warm water                         </td> </tr> <tr> <td style="padding: 5px;"> <b>11</b> • Carry victim into fresh air.                              • Make patient lie down.                              • Give artificial respiration if necessary.                         </td> </tr> <tr> <td style="padding: 5px;"> <b>14</b> • Give water or milk.                              • Then give 4 tablespoons of vegetable oil.                              • Do NOT induce vomiting!                         </td> </tr> </table>	<b>2</b> • Give glass of milk. OR 1 tablespoonful of activated charcoal mixed with a little water. • Induce vomiting (see #1) but not with syrup of ipecac if charcoal was given	<b>5</b> • Give glass of milk, or activated charcoal in water. • Induce vomiting (#1) if not in convulsions. Do not use ipecac if charcoal was given. • Keep patient quiet.	<b>8</b> • Give a glass of milk • Next induce vomiting (#1) • Give 2 teaspoons of bicarbonate of soda in a glass of warm water	<b>11</b> • Carry victim into fresh air. • Make patient lie down. • Give artificial respiration if necessary.	<b>14</b> • Give water or milk. • Then give 4 tablespoons of vegetable oil. • Do NOT induce vomiting!
<b>Keep All Poisons And Medicines Out Of Reach Of Children!</b>	<b>1</b> Induce vomiting with • Finger in throat. OR • 1 tablespoon of syrup of ipecac, followed by a glass of water. OR • Teaspoonful of mustard in half glass of water																			
<b>3</b> • Induce vomiting (See #1) • Then give 4 tablespoons of castor oil • Next give glass of milk OR the white of 2 raw eggs	<b>4</b> • Induce vomiting (see #1) if patient is conscious. • Give glass of milk. OR activated charcoal in water • Give 2 tablespoons of epsom salt in 2 glasses of water. • Keep patient awake																			
<b>6</b> • Induce vomiting (See #1) • Next give 2 tablespoons of epsom salt in 2 glasses of water—except in cases where diarrhea is severe	<b>7</b> • Induce vomiting (See #1) • Give 2 teaspoons of bicarbonate of soda in a glass of warm water • Finally give glass of milk.																			
<b>9</b> • Give activated charcoal in water. • Induce vomiting (see #1) but not with ipecac. • Give 2 tablespoons of epsom salt in 2 glasses of water.	<b>10</b> • Give glass of milk or lime water. • Then induce vomiting. (See #1)																			
<b>12</b> • Induce vomiting (See #1). • Don't let vomit touch victim's skin or yours. • Then give 4 oz mineral oil. Positively do NOT give vegetable or animal oil. • Also give 1 tablespoon of bicarbonate of soda in a quart of warm water	<b>13</b> • Give 2 tablespoons of vinegar in 2 glasses of water • Then give the white of 2 raw eggs... or 2 ounces of vegetable oil. • Do NOT induce vomiting!																			
<b>15</b> • Give patient one or two glasses of milk	<b>16</b> • Give large quantity of water • Give 2 tablespoons of milk of magnesia, or a similar antacid. • Do NOT induce vomiting!																			
<b>2</b> • Give glass of milk. OR 1 tablespoonful of activated charcoal mixed with a little water. • Induce vomiting (see #1) but not with syrup of ipecac if charcoal was given																				
<b>5</b> • Give glass of milk, or activated charcoal in water. • Induce vomiting (#1) if not in convulsions. Do not use ipecac if charcoal was given. • Keep patient quiet.																				
<b>8</b> • Give a glass of milk • Next induce vomiting (#1) • Give 2 teaspoons of bicarbonate of soda in a glass of warm water																				
<b>11</b> • Carry victim into fresh air. • Make patient lie down. • Give artificial respiration if necessary.																				
<b>14</b> • Give water or milk. • Then give 4 tablespoons of vegetable oil. • Do NOT induce vomiting!																				
<b>OVERDOSES</b>  Alcohol • 8 Aspirin • 8 Barbiturates • 9 Belladonna • 5 Bromides • 6 Codeine • 4 Headache & Cold Compounds • 8 Iron Compounds • 7 Morphine, Opium • 4 Paregoric • 4 Pep' Medicines • 2 Sleeping Medicines • 9 Tranquilizers • 9	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; padding: 5px;"> <b>EMERGENCY PHONE NUMBERS</b>                   PHYSICIAN _____                  POISON CONTROL CENTER <u>919-644-8111</u>                  PHARMACIST <u>859-6116</u> </td> <td style="width: 50%; padding: 5px;">                 POLICE _____                  HOSPITAL _____                  AMBULANCE _____                  FIRE _____             </td> </tr> </table>			<b>EMERGENCY PHONE NUMBERS</b>  PHYSICIAN _____ POISON CONTROL CENTER <u>919-644-8111</u> PHARMACIST <u>859-6116</u>	POLICE _____ HOSPITAL _____ AMBULANCE _____ FIRE _____															
<b>EMERGENCY PHONE NUMBERS</b>  PHYSICIAN _____ POISON CONTROL CENTER <u>919-644-8111</u> PHARMACIST <u>859-6116</u>	POLICE _____ HOSPITAL _____ AMBULANCE _____ FIRE _____																			

**Post This Chart**  
**On The Back Of Your**  
**Medicine Cabinet Door**  
**—Or On The Back Of**  
**Your Bathroom Door.**

**TRYON PHARMACY**

620 SOUTH TRADE STREET

PHONE 859-6615

TRYON, N. C. 28782